

# TIPS FOR CONSULTATION

## SAMPLE QUESTIONS

- + What are your qualifications?
- + How much experience do you have in performing this treatment?
- + Which of the products in the Restylane® family is most appropriate for me?
- + What can you do to make me more comfortable during my treatment?
- + How long will the treatment take?
- + What care is recommended after treatment?
- + How much downtime will there be after I receive the treatment (e.g., time away from work, activities, etc.)?
- + When should I schedule my follow-up appointment?
- + What are the most common side effects?
- + What can I do to maintain my results?
- + Are there any important safety considerations that I should be aware of?

**TIP:**  
BE SURE TO CHOOSE A DOCTOR WITH WHOM YOU'RE COMFORTABLE.

## MY QUESTIONS

---

---

---

---

---

---

---

---

## NOTES

---

---

---

---

---

---

---

---



# TIPS FOR CONSULTATION

The Restylane family of products includes Restylane®, Restylane-L®, Restylane Fine Lines-L™, Perlane®, and Perlane-L®. These products are intended for individual use to add volume to facial tissue.

Restylane and Restylane-L are to be used for the correction of wrinkles, lip enhancement and facial contour augmentation.

Perlane and Perlane-L are recommended for shaping the contours of the face, correcting folds and lip enhancement.

Restylane-L, Perlane-L and Restylane Fine Lines-L have been enhanced with the addition of lidocaine in order to reduce discomfort during treatment.

Restylane Fine Lines-L is recommended for the correction of thin, superficial lines, for example the lines around the eyes, the lines around the mouth, the forehead lines and the smile lines.

## IMPORTANT SAFETY CONSIDERATIONS FOR THE RESTYLANE FAMILY OF PRODUCTS

If you have a hypersensitivity to local anesthetics of the amide type, such as lidocaine, you should not use Restylane-L, Perlane-L or Restylane Fine Lines-L.

After your treatment, you might have some swelling, redness, pain, itching, discolouration, and tenderness at the site of treatment. This will typically last less than a week.

Additionally, although rare, red or swollen small bumps may occur in some patients.

In rare circumstances, the doctor may inject into a blood vessel, which can damage the skin, or in very rare cases, cause a temporary loss of vision.

If you are pregnant, breastfeeding or under 18 years, you should not use the Restylane family of products.

The Restylane family of products should not be injected anywhere except the skin or just under the skin.

Like any injection in the skin, the implantation of the Restylane family of products carries a risk of infection.

The use of the Restylane family of products at the site of skin sores, pimples, rashes, hives, cysts or infection should be postponed until healing is complete.

Avoid exposure of the treated area to intense heat (such as solarium and sunbathing) or extreme cold until any initial swelling and redness have resolved.

For each treatment site, a maximum dosage of 0.8 mL of Restylane Fine Lines-L and of 2 mL of Restylane, Restylane-L, Perlane, and Perlane-L per treatment session is recommended.

The Restylane family of products should only be injected by a licensed health care professional. Complete product information and Important Safety Considerations are available at [www.Restylane.ca](http://www.Restylane.ca). Ask your health care professional if the Restylane family of products is an appropriate treatment option for you, as it is not for everyone.

[RESTYLANE.CA](http://RESTYLANE.CA)